# FLAVORED TOBACCO PRODUCTS



Swisher Sweets come in flavors such as grape, strawberry, and chocolate.



Smokeless and spit tobacco are available in flavors such as berry and apple.



E-Cigarette / Vape Liquid, known as e-juice, comes in over 7,000 different flavors, that often come in brightly colored packaging that appeals to youth.<sup>10</sup>

Big Tobacco's own internal documents state sweet products are:

"... for younger people, beginner cigarette smokers, teenagers ... when you feel like a smoke, you want to be reminded of bubble gum."

## WHAT TYPES OF TOBACCO PRODUCTS ARE FLAVORED?

Flavored tobacco products include cigars, chewing tobacco, blunt wraps, electronic cigarettes, and shisha, the tobacco used in hookah. These products help the tobacco industry avoid the Food and Drug Administration's (FDA) 2009 prohibition of flavored cigarettes.¹ Flavored tobacco products are often sold in attractive packaging for a low price. Cigars are frequently available for less than \$1.

### MANY COMMON FLAVORS ARE USED IN TOBACCO PRODUCTS.

Cigars, chewing tobacco, blunt wraps, electronic cigarettes, e-liquid and shisha are sold in fruit, candy, desert and other novelty flavors. Popular flavors include chocolate, pina colada, apple, grape, berry, cotton candy, bubble gum and menthol. The same flavorings used in tobacco products are also used in candy and Kool-Aid.<sup>2</sup> These flavorings are often present in higher amounts in tobacco products than candy.<sup>2</sup>

## WHO USES FLAVORED TOBACCO PRODUCTS?

Flavoring in tobacco products makes it easier for new, young users to tobacco, because the flavoring masks the harshness of the tobacco and enhances the users experience.<sup>3</sup> Nearly 90 percent of addicted adult smokers started smoking by age 18.<sup>4</sup>

Because of their sweet flavors, low cost and attractive packaging, flavored tobacco products are especially appealing to youth. Young people are much more likely to use flavored tobacco products than adults.<sup>3</sup>

According to a report from the Centers for Disease Control and Prevention (CDC), 63 percent of the students who use electronic cigarettes use flavored e-cigarette juice. Also 64 percent of those who use cigars use flavored cigars, and 61 percent of those who use hookah use flavored shisha.<sup>5</sup>

Studies show that young people perceive flavored tobacco products as tasting better and being safer than unflavored products, even though they are just as dangerous and addictive as cigarettes.

## WHAT DOES THE TOBACCO INDUSTRY HAVE TO SAY ABOUT FLAVORED TOBACCO PRODUCTS?

Documents from the tobacco settlements revealed the "Graduation Theory," a method used by the tobacco industry aimed to secure customer loyalty.8 This approach implies that new users start with milder tasting and flavored products. Then, they graduate to full-bodied, less flavored items that often contain more nicotine, resulting in a life-long addiction.







Shisha is the tobacco used in hookahs. It is often flavored.



Establishment: Red Carpet Location: 2921 North 11th Street, Bismarck, ND 58503 Across the street from Century High School Date Photo was Taken: 3/17/2015



## WHAT CAN COMMUNITIES DO ABOUT FLAVORED TOBACCO PRODUCTS?

Flavored tobacco products are a major public health concern because they encourage young people to start using tobacco. While the FDA banned flavored cigarettes other than menthol in 2009, the ban does not affect other tobacco products, many of which are now heavily marketed by the tobacco industry. The use of these products especially among young people, has spiked.<sup>5</sup>

State and local governments can adopt laws that restrict the sale of flavored tobacco products within their jurisdiction. Limiting the sale of products, setting a minimum price or creating a minimum pack size for cigars are some of the ways communities can protect their youth. According to the 2000 U.S. Surgeon Generals Report, raising prices of tobacco products decreases tobacco use among youth and young adults? Making flavored tobacco products less accessible also has the potential to reduce youth tobacco use.

# WHAT COMMUNITIES HAVE TAKEN A STANCE AGAINST FLAVORED TOBACCO PRODUCTS?

Locally, Minneapolis, Saint Paul, and Shoreview restricted the sale of flavored tobacco, allowing flavored tobacco to only be sold in adult-only tobacco shops, meaning convenience stores can no longer sell flavored tobacco products. This significantly reduces availability and visibility of the products.

New York City and Providence, R.I., passed policies restricting flavored tobacco sales, with the exception of menthol-flavored products. Chicago passed an ordinance restricting sale of all flavored products, including menthol cigarettes, within 500 feet of schools except in adult-only tobacco retailers.

#### **SOURCES**

- 1 One Hundred Eleventh United States Congress. (2009). Family Smoking Prevention and Tobacco Control Act. Retrieved from <a href="https://www.qpo.gov/fidsys/pkg/PLAW-111publ31/htm//PLAW-111publ31.htm">https://www.qpo.gov/fidsys/pkg/PLAW-111publ31/htm//PLAW-111publ31.htm</a>
- 2 Brown, J. E., Kuo, W., Is-abelle, L. M., & Pankow, J. F. (2014, June 5). Candy Flavorings in Tobacco. Retrieved from <a href="http://www.nejm.org/doi/full/10.1056/NEJMc1403015#">http://www.nejm.org/doi/full/10.1056/NEJMc1403015#</a>=article
- 3 King, B. A., Dube, S. R., & Tynan, M. A. (2013, February). Flavored Cigar Smoking Among U.S. Adults: Findings From the 2009-2010 National Adult Tobacco Survey. Nicotine & Tobacco Research, 15(2), 608-614. doi:10.1093/ntr/nts178
- 4 U.S. Department of Health and Human Services. (2012). Preventing Tobacco Use Among Youth and Young Adults: A Report of the Surgeon General. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. Retrieved from
- http://www.surgeongeneral.gov/library/reports/preventing-youth-tobacco-use/full-report.pdf
- 5 U.S. Centers for Disease Control & Prevention, (2015, October). Flavored Tobacco Product Use Among Middle and High School Students -United States, 2014. Morbidity and Mortality Weekly Report. Retrieved from <a href="http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6438a2.htm?scid=mm6438a2">https://www.cdc.gov/mmwr/preview/mmwrhtml/mm6438a2.htm?scid=mm6438a2</a> w
- 6 U.S. Food and Drug Administration. (n.d.). Flavored Tobacco Product Factsheet. Retrieved from <a href="http://www.fda.gov/TobaccoProducts/Labeling/ProductsIngredientsComponents/ucm183198.htm#4">http://www.fda.gov/TobaccoProducts/Labeling/ProductsIngredientsComponents/ucm183198.htm#4</a>
- 7 Report from R.M. Manko Assoc. to Lorillard Tobacco Co. (1978, August)
- 8 David Weiss Association. (1984, November 16). The Graduation Theory. Retrieved from <a href="http://legacy.library.ucsf.edu/tid/lfc46b00/pdf?search=%22graduation%20theory%22">http://legacy.library.ucsf.edu/tid/lfc46b00/pdf?search=%22graduation%20theory%22</a>. Bates No. USSTC1945141-USSTC1945142
- 9 U.S. Department of Health and Human Services. (2000). Reducing Tobacco Use: A Report of the Surgeon General. Retrieved from <a href="http://www.surgeongeneral.gov/library/reports/preventing-youth-tobacco-use/full-report.pdf">http://www.surgeongeneral.gov/library/reports/preventing-youth-tobacco-use/full-report.pdf</a>
- 10 New England Journal of Medicine: Nicotine Poisoning in an Infant, N Engl J Med 2014; 3070-2249-2250 September 6, 2017 DOI: 10.1056/nejmC1403843